

Rapid Analysis of Amino Acids Using the Hitachi LaChromUltra® **High Speed Liquid Chromatography System**

Hitachi High Technologies America, Inc.

mino acid analysis is an important technique involved in a wide range of bioanalytical applications including protein and peptide analysis, monitoring of amino acid uptake in cell culture media, quality assessment of foodstuffs and clinical diagnosis of Historically, this technique has been metabolic disorders. performed on highly specialized chromatographic systems dedicated for comprehensive (up to 50-plus components) and quantitative amino acid analysis. While the data from these platforms are extremely reproducible and informative, analysis time can be lengthy (up to 150 minutes) and some sensitivity may be lost. Here we describe an alternative methodology for the rapid analysis of amino acids suitable for the analysis of proteins and peptides, cell culture media, and foodstuffs using the highly flexible Hitachi LaChromUltra® U-HPLC system. pre-column derivatization using 4-fluoro-7-nitrobenzo-furazan (NBD-F), amino acids are analyzed via reversed phase separation and ultraviolet or fluorescence detection. The data here specifically describe the analysis of a commercial sports drink via fluorescence detection in under eight minutes¹.


Experimental Conditions

Experimental Conditions		
Module	Conditions	
Pump (L-2160U)	Mobile Phase A: 120mM Ammonium Formate + 1g/L Tetrabutylammonium Bromide/H3PO4 concentration (v/v): 0.15% Mobile Phase B: Acetonitrile/Tetrahydrofuran (95:5) + 1g/L Tetrabutylammonium Bromide Flow Rate: 0.55mL/min	
Autosampler (L-2200U)	Injection Volume: 10μL	
Oven (L-2300)	Temperature: 37°C	
Detector (L-2485U)	Ex: 470nm; Em: 540nm	
Standard	23 amino acid mixture	


Results – Performance

Peak Area Reproducibility	0.38 – 0.95% RSD
Linearity (0.2 – 20.0pmol)	0.9991 – 0.9999% RSD
Detection Limit (S/N=3)	<40fmol/10uL (except Pro=60, Orn=70, Cys2=110 and Met=120)
Quantitative Limit (S/N=10)	<100fmol/10uL (except Pro=200, Orn=230, Cys2=375 and Met=400)

Results – Chromatogram of 23-Component Mixture



Results – Chromatogram of Commercial Sports Drink



Discussion

Hitachi's LaChromUltra® liquid chromatography system is extremely effective at rapid analysis of amino acids in a variety of matrices. The system exhibits a linear response over a broad concentration range, excellent reproducibility and very low detection and quantitation limits.

Reference:

1 - Technical Data, Hitachi High Technologies Corporation.

Hitachi High Technologies America, Inc.

Life Sciences Division 5100 Franklin Drive Pleasanton, CA 94588 Toll Free: (800) 548-9001